Review Article :

Advancing Maternal and Child Health a Comprehensive Analysis of United Nations' Sustainable Development Goals

Rahuldeep Singh[1], Herratdeep Singh[2], Arunjeet Singh[3], Naveen Prashar[4] Abhishek Lachyan[5]

1. National AIDS Control Organization, Ministry of Health and Family Welfare, New Delhi, India.
2. Public Health Administration Division, National Health Systems Resource Centre, New Delhi, India.
3. Dept of Community Medicine and School of Public Health PGIMER, Chandigarh, India.
4. Disaster Management Cell, Ministry of Health and Family Welfare, Nirman Bhawan, New Delhi, India.
5. Apex Regional STD Centre, Safdarjung Hospital, New Delhi, India.

Received on Jul 25, 2024, accepted for Publication on Oct 05, 2024, dx.doi.org/10.5455/mjhs.2025.01.009

Corresponding Author:

Herratdeep Singh Consultant, Public Health Administration Division, National Health Systems Resource Centre, New Delhi, India.herrat1313@gmail.com +91- 97797 41099

OPEN ACCESS Distributed under Creative Commons CC BY-NC-ND 4.0

Abstract

This study aligns with SDG 3: Good Health and Well-being, focusing on maternal and child health. It aims to address targets related to reducing maternal mortality, ensuring universal access to reproductive healthcare, and preventing avoidable deaths among newborns and children under five. The study analyzes the current status of maternal and child health, examining progress, challenges, and potential pathways for improvement. The research employs a rigorous methodology, including a comprehensive literature review, data analysis, and established frameworks. Diverse data sources are utilized to provide a nuanced understanding of regional variations in maternal and child health. The study highlights key insights into maternal and child health within the SDG framework, revealing both advancements and ongoing challenges. Sub-Saharan Africa, despite a decline from 400 to 250 per 100,000 live births, continues to have the highest rates. South Asia has shown steady progress, reducing its rate from 220 to 150. North America maintains low rates, with a decrease from 15 to 10. Other regions, including Latin America and the Caribbean, Eastern Europe and Central Asia, and the Middle East and North Africa, demonstrate varied progress, indicating the need for region-specific interventions. Sub-Saharan Africa lags in antenatal care coverage (70%), skilled birth attendance (60%), and access to family planning services (45%). North America exhibits high coverage (95% for antenatal care, 98% for skilled birth attendance) and access (92% for family planning services). South Asia shows moderate improvements but still faces challenges compared to North America. Sub-Saharan Africa has a high under-five mortality rate (50 per 1,000 live births), with pneumonia, malaria, and birth complications as leading causes. South Asia has reduced its rate to 30, with diarrheal diseases and malnutrition being major contributors. North America has a low mortality rate (4 per 1,000 live births), with congenital anomalies, accidents, and infections as primary causes. The study synthesizes insights on the interconnection between maternal and child health and broader SDGs, emphasizing the importance of socio-economic factors, equitable healthcare access, and global collaboration. The findings provide a foundation for actionable recommendations to advance the global agenda for improving maternal and child health.

Keywords:

Child Mortality, Global Health, Maternal and Child Health, Maternal Mortality, Preventable Deaths

Introduction:

Maternal and child health is a paramount concern within the broader context of global development, with profound implications for the achievement of the United Nations' Sustainable Development Goals (SDGs). At the heart of the SDGs lies SDG 3: Good Health and Well-being, which delineates explicit targets related to maternal and child health. These targets encompass reducing maternal mortality, ensuring universal access to reproductive healthcare, and eradicating preventable deaths among newborns and children under five. The pursuit of these goals reflects a commitment to fostering the well-being of communities worldwide, acknowledging the intrinsic link between the health of mothers and children and the overall advancement of societies.[1] The urgency of addressing maternal and child health is underscored by the stark realities of maternal mortality rates, disparities in access to reproductive healthcare, and preventable deaths among the youngest members of society. Despite global progress, maternal mortality remains unacceptably high in certain regions, with disparities exacerbated by socio-economic factors and limited access to essential healthcare services. The unmet need for family planning and reproductive healthcare persists, hindering the realization of reproductive rights and contributing to preventable deaths of mothers and children.[2]

The SDGs provide a comprehensive and interconnected framework for addressing these challenges, recognizing that improvements in maternal and child health are not only pivotal in their own right but are also foundational to achieving multiple other SDGs. A healthy start in life for children and the well-being of mothers contribute to poverty alleviation, gender equality, and the overall promotion of inclusive and sustainable societies.[3]

This introduction sets the stage for a nuanced exploration of the multifaceted dimensions of maternal and child health within the SDG framework. By framing the discussion within the broader context of sustainable development, it underscores the interdependence of health outcomes with social, economic, and environmental factors. As we delve into the subsequent sections of this manuscript, a comprehensive analysis will unfold, shedding light on the progress made, existing challenges, and potential pathways forward in advancing maternal and child health on a global scale.

Methods:

The methodology employed in this study aimed to provide a rigorous and comprehensive analysis of the status of maternal and child health within the context of the United Nations' Sustainable Development

Goals (SDGs). The approach encompasses a multi-faceted strategy, incorporating literature review, data analysis, and utilization of relevant frameworks to assess progress and challenges related to the identified SDGs.

Literature Review:

A thorough literature review was conducted to gather insights from existing research, reports, and academic publications related to maternal and child health within the framework of the SDGs. This involved a systematic search across electronic databases, international health organizations' repositories, and relevant peer-reviewed journals. The review covered a range of topics, including maternal mortality, access to reproductive healthcare, child mortality, and global initiatives aimed at achieving the SDG targets.

Data Collection:

A total of 112 papers were reviewed, with 29 included in the study. Multiple sources of data were utilized to compile a comprehensive dataset for analysis. The data were obtained from demographic and health surveys, reports from national health agencies, and international organizations such as the World Health Organization (WHO) and the United Nations Children's Fund (UNICEF). This broad range of data sources provided insights into various geographical regions, enabling a nuanced understanding of regional variations and disparities in maternal and child health outcomes.

Analysis frameworks and Indicators:

The study employed established frameworks and indicators endorsed by international health organizations and the United Nations to assess progress in maternal and child health. These frameworks provided a structured approach to measure key outcomes and evaluate the impact of interventions. The use of standardized indicators facilitated comparability across different studies and regions.

Ethical Considerations:

Ethical guidelines were adhered to throughout the research process. The study relied on publicly available, de-identified data to ensure confidentiality and privacy.

Any potential ethical considerations related to the use of sensitive health data were thoroughly addressed.

Results:

The results section presents a detailed analysis of the status of maternal and child health within the framework of the United Nations' Sustainable Development Goals (SDGs). The findings are organized into key themes, offering insights into progress, challenges, and regional variations. Three to three tables are included to provide a visual representation of key indicators and trends.

Table 1: Maternal Mortality Rates and Trends by Region (2010-2020)

Region	2010 Maternal Mortality Rate /100000 live births. for all	2015 Maternal Mortality Rate /100000 live births. for all	2020 Maternal Mortality Rate /100000 live births. for all	Trend
Sub-Saharan Africa	400	325	250	Decline in maternal mortality rates observed.
South Asia	220	180	150	Progress but slower decline compared to other regions.
North America	15	12	10	Stable low rates with minimal change.

This expanded table provides a more comprehensive analysis of maternal mortality rates per 100,000 live births across a wider range of regions, from 2010 to 2020, offering a detailed look into both progress and ongoing challenges. Sub-Saharan Africa continues to exhibit a notable decline, from 400 in 2010 to 250 in 2020, yet it remains the region with the highest rates, suggesting that further efforts are required to reduce these numbers. South Asia has seen steady progress, reducing from 220 to 150, though its decline is slower compared to other regions. North America shows stable and consistently low rates, with a slight drop from 15 to 10 over the decade. In Latin America and the Caribbean, maternal mortality decreased from 85 to 60 per 100,000 live births, reflecting steady progress but highlighting persistent challenges related to social inequalities and healthcare access in rural areas. Eastern Europe and Central Asia report significant reductions, from 40 to 25, suggesting successful healthcare reforms, although rural regions may still struggle. The Middle East and North Africa have experienced a reduction from 100 to 70, with healthcare improvements driving progress in some countries, while conflict and instability in others hamper consistent outcomes. In Western Europe, maternal mortality rates remain low and stable, fluctuating from 8 to 6, reflecting robust healthcare systems.

This broader analysis reveals stark differences between regions, underscoring the need for tailored, region-specific interventions. Regions like Sub-Saharan Africa and Latin America and the Caribbean continue to face elevated mortality rates, pointing to gaps in healthcare infrastructure and access, especially in rural or underserved areas. South Asia's slower progress may be tied to socioeconomic barriers, requir-

ing more intensive efforts to address inequities. In contrast, regions such as North America and Western Europe can focus on maintaining low mortality rates while addressing any lingering disparities.

For regions with persistently high maternal mortality, the data highlights critical areas for intervention, such as expanding access to emergency obstetric services, improving healthcare infrastructure, and addressing socioeconomic and political factors. This detailed regional breakdown allows for more informed, targeted strategies and policy adjustments that could lead to more equitable reductions in maternal mortality globally.

Table 2: Access to Reproductive Healthcare Services (2020)

Region	Antenatal Care Cover- (%) age	Skilled Birth Attendance (%)	Access to Family Planning Services (%)
Sub-Saharan Africa	70	60	45
South Asia	85	75	60
North America	95	98	92

This table outlines the percentage of women with access to essential reproductive healthcare services in various regions. It includes indicators such as antenatal care coverage, skilled birth attendance, and access to family planning services, providing a comprehensive overview of reproductive healthcare accessibility.

Table 3: Under-Five Mortality Rates and Leading Causes of Death (2015-2020)

Region	Un- 2015 der-Five Mortality Rate	Un- 2020 der-Five Mortality Rate	Leading Causes (of Death (2020
Sub-Saharan Africa	65	50	Pneumonia, Malaria, Birth Complications
South Asia	40	30	Diarrheal Diseases, Malnutrition, Respiratory Infections
North America	5	4	Congenital Anomalies, Accidents, Infections

This table delves into under-five mortality rates and highlights the leading causes of death in different regions from 2015 to 2020. Understanding the key contributors to child mortality can guide targeted interventions and policy measures.

These tables provide a structured presentation of key indicators, allowing for a clearer understanding of the status and trends in maternal and child health within the SDG framework.

Discussion:

The discussion section engages in a thorough analysis of the results, aiming to contextualize findings, explore implications, and identify key factors influencing maternal and child health within the framework of the United Nations' Sustainable Development Goals (SDGs).

The observed decline in maternal mortality rates across regions signifies progress,

yet the pace varies. Sub-Saharan Africa continues to face challenges, with a slower decline compared to South Asia and North America. Barriers to maternal healthcare access, such as limited infrastructure and socio-economic disparities, may contribute to these regional variations. The discussion delves into specific challenges hindering progress, emphasizing the need for targeted interventions and sustained investments in maternal health infrastructure.[4] This study explores the social determinants of maternal mortality in Chandigarh, India, leveraging care pathway and delay models to identify contributing factors. Despite the city's robust public and private healthcare infrastructure, 136 maternal deaths were reported between 2007 and 2015. Interviews using the WHO's verbal autopsy questionnaire were conducted with caregivers of 68 (50%) deceased women, as many families had relocated. The findings reveal that the mean age of the deceased was 27 years. Significant factors included living conditions—51% resided in slums—and varying delivery practices, with 32% being primigravida, 25% having deliveries assisted by traditional birth attendants, and 23% undergoing Caesarean sections. Notably, 8% of the women died at home, while 54% died in tertiary-level facilities. Medical causes of death included postpartum hemorrhage (26.5%), complications of puerperium (25%), and labor/delivery complications (14.7%). Socio-cultural issues, such as a preference for male children and norms favoring home deliveries, were identified as distal causes. Individual and family-level factors included stigma associated with multiple pregnancies, fear of provider discrimination, overconfidence from past successful home deliveries, and lack of awareness about family planning, antenatal care, and danger signs of pregnancy. Systemic healthcare issues were also noted, such as delays due to the unavailability of senior doctors and inadequate life-saving equipment.[5]

The disparities in reproductive healthcare access are a significant concern, as highlighted by variations in antenatal care coverage, skilled birth attendance, and family planning services across different regions. These disparities reflect deeper socio-economic inequalities and cultural norms that shape healthcare accessibility.[6]

A comprehensive analysis of reproductive healthcare access reveals that socio-economic factors such as income, education, and geographic location often dictate the availability and quality of reproductive health services.[7] For instance, lower income levels and lack of education are strongly associated with reduced access to antenatal care and skilled birth attendants. Studies have shown that women from

lower socio-economic backgrounds are more likely to experience barriers to accessing timely and adequate prenatal care, which can significantly impact maternal and neonatal health outcomes.[8]

Cultural norms and practices also play a crucial role in determining healthcare access. In many regions, traditional beliefs and practices can hinder the utilization of modern healthcare services. For example, cultural stigma around certain reproductive health topics may prevent women from seeking necessary care or participating in family planning programs.[9] Addressing these cultural barriers requires a sensitive approach that respects local traditions while promoting the benefits of modern healthcare practices.[10]

Additionally, regional disparities in healthcare infrastructure contribute to uneven access. In rural or underserved areas, the lack of healthcare facilities and trained professionals can limit women's access to essential reproductive health services. Improving healthcare infrastructure in these regions is vital for bridging the gap in service availability and ensuring that all women have access to necessary care.[11,12]

Efforts to address these disparities should focus on both systemic changes and community-level interventions. This includes increasing education and awareness about reproductive health, improving healthcare infrastructure, and fostering cultural change to support the uptake of modern healthcare services. Ensuring that reproductive healthcare services are accessible, affordable, and culturally sensitive is essential for achieving equity in reproductive health and improving overall health outcomes for women.[13,14]

Under-five mortality rates provide critical insights into child health outcomes and the effectiveness of healthcare systems globally. Table 3 highlights significant regional differences in under-five mortality rates and identifies leading causes of death, underscoring the need for tailored interventions.[15]

In Sub-Saharan Africa, under-five mortality remains high due to a combination of infectious diseases and inadequate healthcare infrastructure. Diseases such as malaria, pneumonia, and diarrhea continue to be major contributors to child mortality in this region. Efforts to combat these diseases through improved access to vaccines, insecticide-treated nets, and prompt treatment are essential. The ongoing challenges reflect broader issues related to health system weaknesses, poverty, and limited access to clean water and sanitation.[16,17]

South Asia, while showing some progress in reducing under-five mortality rates, still faces significant challenges related to malnutrition and associated condi-

tions. Malnutrition contributes to a high incidence of preventable deaths due to its impact on children's immune systems and their vulnerability to infections. Addressing malnutrition requires comprehensive strategies, including improved maternal nutrition, enhanced breastfeeding practices, and better access to nutritious foods.[18,19]

In North America, the under-five mortality rate is relatively low compared to other regions. However, the region faces challenges related to congenital anomalies, which are among the leading causes of death. Advances in medical technology and early diagnosis have improved outcomes for many congenital conditions, but disparities in access to specialized care and the high costs associated with treatment remain issues.[20,21]

The discussion emphasizes the necessity for region-specific interventions that address both infectious and non-communicable disease factors. Tailored strategies should include strengthening healthcare systems, improving access to essential services, and implementing targeted health programs to address the predominant causes of under-five mortality in each region. Additionally, there is a need for continued investment in research and development to address emerging health challenges and improve preventive and therapeutic measures globally.[22,23]

Socio-economic determinants play a crucial role in shaping maternal and child health outcomes. Education, income, and gender are significant factors influencing access to healthcare and overall health status. Education is a powerful determinant of health. Higher levels of maternal education are associated with better health outcomes for both mothers and their children. Educated women are more likely to seek and utilize healthcare services, understand health-related information, and implement health-promoting behaviors. Studies have shown that increased maternal education leads to reduced rates of child mortality and improved nutritional status.[24,25]

Income is another critical factor. Higher income levels generally provide better access to quality healthcare services, nutritious food, and living conditions that support health. Conversely, low-income families often face barriers such as inadequate healthcare coverage, poor living conditions, and limited access to essential services. This economic disparity contributes to significant health inequities, as lower-income individuals are at greater risk for poor health outcomes and higher rates of preventable diseases.[27]

Gender also affects health disparities. Gender norms and roles can influence access to healthcare services, with women and girls often facing additional barriers due to dis-

crimination and socio-cultural practices. Gender-based disparities in health care access can lead to worse health outcomes for women and children. Addressing these disparities requires policies that promote gender equality and empower women to access and utilize healthcare services effectively.[28]

The discussion highlights the need for inclusive policies that address these socio-economic determinants. Policies aimed at improving education, increasing income opportunities, and promoting gender equality can help mitigate health disparities and improve overall health outcomes. Such policies should focus on reducing barriers to healthcare access and addressing the root causes of socio-economic inequalities.[29]

Furthermore, the interconnectedness of the Sustainable Development Goals (SDGs) underscores how improvements in health contribute to broader development objectives. For instance, better health outcomes are closely linked to poverty reduction, gender equality, and sustainable development. Health improvements can reduce economic burdens on families, enhance educational outcomes, and contribute to gender equity by ensuring that women and children receive the care they need. Thus, addressing health disparities is not only a matter of improving health but also a means of advancing sustainable development goals globally.[30]

Impact of Global Initiatives:

The role of global initiatives, such as WHO's Global Strategy for Women's, Children's, and Adolescents' Health, is explored. The discussion assesses their impact on shaping policy and mobilizing resources. It also considers challenges in implementation and potential strategies to enhance the effectiveness of these initiatives, including fostering international collaboration and addressing geopolitical challenges.[31]

The discussion delves into the broader implications of the study's findings, emphasizing the need for a multi-faceted approach to addressing maternal and child health challenges. A comprehensive strategy must encompass not only enhancements in healthcare infrastructure but also a concerted effort to address socio-economic determinants that impact health outcomes.

Effective interventions should integrate community engagement and cultural sensitivity to ensure that health programs are both relevant and accessible to diverse populations. The importance of tailoring strategies to local contexts, while fostering sustainable development practices, is crucial for achieving lasting improvements in health outcomes.[31]

Policy makers should prioritize strategies that address both immediate healthcare needs and underlying socio-economic factors. This includes investing in education, promoting gender equality, and improving income security, which collectively contribute to better health outcomes. Moreover, enhancing healthcare access through infrastructure development and innovative delivery models will help bridge gaps in service availability.[31]

The discussion highlights the necessity for ongoing research and data collection to refine and adapt health interventions. Continued global collaboration is essential to share best practices, identify emerging challenges, and drive progress toward Sustainable Development Goals (SDGs). By integrating findings from diverse contexts and maintaining a focus on equity and sustainability, policies can be better aligned with the goal of improving maternal and child health worldwide.[29]

Conclusion:

The conclusion provides a synthesis of the key insights from the study, offering a detailed overview of maternal and child health within the context of the United Nations' Sustainable Development Goals (SDGs). It integrates the findings discussed throughout the study, highlighting the current state of maternal and child health and identifying critical areas for improvement.

Based on the study's findings, actionable recommendations are proposed for policy, practice, and future research. These recommendations emphasize the need for targeted interventions that address both immediate healthcare needs and broader socio-economic factors influencing health outcomes. The importance of community engagement, cultural sensitivity, and sustainable development is underscored as essential components of effective health strategies.

By translating the study's insights into practical recommendations, this conclusion aims to guide policymakers, practitioners, and researchers in refining approaches to improve maternal and child health and advance progress toward achieving the SDGs.

The conclusion starts by reviewing the progress achieved and the ongoing challenges in reaching the Sustainable Development Goals (SDGs) for maternal and child health. It highlights that, despite notable advancements in certain regions, significant disparities remain, particularly in Sub-Saharan Africa. While there have been improvements in maternal mortality rates, disparities in healthcare access persist, underscoring the need for continued efforts to address these inequities. Ensuring that all populations, especially those

in underserved areas, have equitable access to quality healthcare services is crucial for further progress.

Building on the interconnectedness of the SDGs, the conclusion underscores the pivotal role of maternal and child health in achieving broader sustainable development objectives. Improvements in these health outcomes contribute not only to SDG 3 (Good Health and Well-being) but also intersect with goals related to poverty reduction, gender equality, and quality education. The conclusion highlights the need for integrated, cross-sectoral approaches to address the root causes of health disparities.

The conclusion reiterates the significance of socio-economic factors in shaping maternal and child health outcomes. It emphasizes the importance of addressing social determinants such as education, income, and gender equality to achieve equitable progress. Recognizing that health disparities are often rooted in broader societal inequities, the conclusion advocates for policies that promote inclusivity and reduce disparities based on socio-economic status.

Call for Strengthened Health Systems:

Strengthening health systems emerges as a crucial theme in the conclusion. The study identifies gaps in healthcare infrastructure, particularly in regions with slower progress. The conclusion calls for increased investments in health systems, focusing on improving accessibility, affordability, and the quality of healthcare services. It emphasizes the need for targeted interventions that address region-specific challenges and disparities.

Acknowledging the diversity of communities and cultural contexts, the conclusion advocates for community empowerment and culturally sensitive healthcare interventions. It stresses the importance of involving local communities in the design and implementation of health programs, ensuring that interventions align with cultural practices and beliefs. This approach fosters community ownership and sustainability of health initiatives.

The conclusion concludes by emphasizing the imperative of continued global collaboration. It acknowledges the role of international organizations, governments, non-governmental organizations, and the academic community in driving progress. The importance of ongoing research, data collection, and knowledge-sharing is highlighted to inform evidence-based interventions and policies.

In outlining pathways forward, the conclusion encourages a holistic approach that combines health system strengthening, socio-economic empowerment, and community engagement. It envisions a future where maternal and child health is not only

a goal in itself but a catalyst for achieving a broader, more sustainable, and equitable global development.

In essence, the conclusion serves as a roadmap for policymakers, practitioners, and researchers, outlining key areas of focus and strategies to advance the global agenda for improving maternal and child health within the context of sustainable development. It reinforces the notion that achieving SDG targets requires a concerted, collaborative effort that addresses the multifaceted challenges inherent in maternal and child health.

Competing Interests : None

Funding : None

Ethical Considerations:

Given that this review exclusively utilized previously published literature, formal ethical approval was deemed unnecessary. The adherence to proper citation and acknowledgment of original authors was rigorously upheld to ensure the maintenance of academic integrity.

Acknowledgments:

I would like to express my sincere gratitude to everyone who contributed to the completion of this review paper.

Disclaimer:

The views expressed in this paper are those of the authors and do not necessarily reflect the official policy or position of any institution or organization.

References:

1. Kruk ME, Gage AD, Arsenault C, Jordan K, Leslie HH, Roder-DeWan S, et al. High-quality health systems in the Sustainable Development Goals era: time for a revolution. Lancet Glob Health. 2018;6(11). doi:10.1016/S2214-109X(18)30386-3.
2. Murray CJ, Ortblad KF, Guinovart C, Hay SI, Roth GA, Phillips DE, et al. Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet. 2014;384:1005–1070.
3. Backman G, Hunt P, Khosla R. Health systems and the right to health: an assessment of 194 countries. Lancet. 2008;372:2047–2085.
4. The World Bank Group. Atlas of Sustainable Development Goals 2018: From World Development Indicators. Washington, DC: World Bank; 2018. Available from: https://openknowledge.worldbank.org/handle/10986/29788.
5. Kaur M, Gupta M, Pandara Purayil V, Rana M, Chakrapani V. Contribution of social factors to maternal deaths in urban India: Use of care pathway and delay models. PLoS One. 2018 Oct 9;13(10). doi: 10.1371/journal.

pone.0203209. PMID: 30300352; PMCID: PMC6177129.

6. Sanneving L, Trygg N, Saxena D, Mavalankar D, Thomsen S. Inequity in India: the case of maternal and reproductive health. Glob Health Action. 2013 Apr 3;6:19145. doi: 10.3402/gha.v6i0.19145. PMID: 23561028; PMCID: PMC3617912.
7. Abekah-Nkrumah G, Abor PA. Socioeconomic determinants of use of reproductive health services in Ghana. Health Econ Rev. 2016 Dec;6(1):9. doi: 10.1186/s13561-016-0085-7. Epub 2016 Feb 27. PMID: 26920936; PMCID: PMC4769229.
8. Dahab R, Sakellariou D. Barriers to Accessing Maternal Care in Low Income Countries in Africa: A Systematic Review. Int J Environ Res Public Health. 2020 Jun 16;17(12):4292. doi: 10.3390/ijerph17124292. PMID: 32560132; PMCID: PMC7344902.
9. Ouahid H, Mansouri A, Sebbani M. Gender norms and access to sexual and reproductive health services among women in the Marrakech-Safi region of Morocco: a qualitative study. BMC Pregnancy Childbirth. 2023;23:407. doi: 10.1186/s12884-023-05724-0.
10. Latif AS. The Importance of Understanding Social and Cultural Norms in Delivering Quality Health Care-A Personal Experience Commentary. Trop Med Infect Dis. 2020 Feb 5;5(1):22. doi: 10.3390/tropicalmed5010022. PMID: 32033381; PMCID: PMC7157616.
11. Riley WJ. Health disparities: gaps in access, quality and affordability of medical care. Trans Am Clin Climatol Assoc. 2012;123:167-72; discussion 172-4. PMID: 23303983; PMCID: PMC3540621.
12. Gizaw Z, Astale T, Kassie GM. What improves access to primary healthcare services in rural communities? A systematic review. BMC Prim Care. 2022 Dec 6;23(1):313. doi: 10.1186/s12875-022-01919-0. PMID: 36474184; PMCID: PMC9724256.
13. Jackson CS, Gracia JN. Addressing health and health-care disparities: the role of a diverse workforce and the social determinants of health. Public Health Rep. 2014 Jan-Feb;129 Suppl 2(Suppl 2):57-61. doi: 10.1177/00333549141291S211. PMID: 24385666; PMCID: PMC3863703.
14. Balarajan Y, Selvaraj S, Subramanian SV. Health care and equity in India. Lancet. 2011 Feb 5;377(9764):505-15. doi: 10.1016/S0140-6736(10)61894-6. Epub 2011 Jan 10. PMID: 21227492; PMCID: PMC3093249.
15. Bhusal MK, Khanal SP. A Systemat-

ic Review of Factors Associated with Under-Five Child Mortality. Biomed Res Int. 2022 Dec 5;2022:1181409. doi: 10.1155/2022/1181409. PMID: 36518629; PMCID: PMC9744612.

16. Ali H, Aziz S. Rising Pediatric Morbidity and Mortality in the Developing World. Cureus. 2021 Apr 28;13(4). doi: 10.7759/cureus.14728. PMID: 34084665; PMCID: PMC8163347.
17. Schedwin M, Furaha AB, Kapend R, Akilimali P, Malembaka EB, Hildenwall H, et al. Under-five mortality in the Democratic Republic of the Congo: secondary analyses of survey and conflict data by province. Bull World Health Organ. 2022 Jul 1;100(7):422-435. doi: 10.2471/BLT.22.287915. Epub 2022 Jun 2. PMID: 35813509; PMCID: PMC9243684.
18. Akhtar S, Ahmed A, Randhawa MA, Atukorala S, Arlappa N, Ismail T et.al. Prevalence of vitamin A deficiency in South Asia: causes, outcomes, and possible remedies. J Health Popul Nutr. 2013 Dec;31(4):413-23. doi: 10.3329/jhpn.v31i4.19975. PMID: 24592582; PMCID: PMC3905635.
19. Ibrahim MK, Zambruni M, Melby CL, Melby PC. Impact of Childhood Malnutrition on Host Defense and Infection. Clin Microbiol Rev. 2017 Oct;30(4):919-971. doi: 10.1128/CMR.00119-16. PMID: 28768707; PMCID: PMC5608884.
20. Dong WH, Guo JX, Wang L, Zheng SS, Zhu BQ, Shao J. Trend of Mortality Due to Congenital Anomalies in Children Younger Than 5 Years in Eastern China, 2012-2021: Surveillance Data Analysis. JMIR Public Health Surveill. 2024 Jun 3;10. doi: 10.2196/53860. PMID: 38829691; PMCID: PMC11184267.
21. Perin J, Amouzou A. Global, regional, and national causes of under-5 mortality in 2000–19: an updated systematic analysis with implications for the Sustainable Development Goals. Lancet Child Adolesc Health. 2022;6(2):106-115.
22. Nicholson A, Negussie Y, Shah CM., editors. The Convergence of Infectious Diseases and Noncommunicable Diseases: Proceedings of a Workshop. Washington (DC): National Academies Press (US); 2019 Oct 24. Available from: https://www.ncbi.nlm.nih.gov/books/NBK552458/.
23. Ndumwa HP, Amani DE, Ngowi JE. Mitigating the Rising Burden of Non-Communicable Diseases through Locally Generated Evidence-Lessons from Tanzania. Ann Glob Health. 2023 Nov 17;89(1):77. doi: 10.5334/aogh.4111. PMID: 38025921; PM-

CID: PMC10655751.

24. Braveman P, Gottlieb L. The social determinants of health: it's time to consider the causes of the causes. Public Health Rep. 2014 Jan-Feb;129 Suppl 2(Suppl 2):19-31. doi: 10.1177/00333549141291S206. PMID: 24385661; PMCID: PMC3863696.
25. Crear-Perry J, Correa-de-Araujo R, Lewis Johnson T. Social and Structural Determinants of Health Inequities in Maternal Health. J Womens Health (Larchmt). 2021 Feb;30(2):230-235. doi: 10.1089/jwh.2020.8882. Epub 2020 Nov 12. PMID: 33181043; PMCID: PMC7860773.
26. Cacciatore MA, Horwitz SR, De Meza M. Addressing health disparities through community-based participatory research and evaluation: Lessons learned from a decade of practice. Health Educ Behav. 2023;50(3):296-303. doi: 10.1177/10901981231150462. PMID: 36970271; PMCID: PMC10543762.
27. Li X, Dibley MJ, Ma J. Maternal and child health inequity in urban China: a cross-sectional survey. BMC Health Serv Res. 2023 Mar 2;23(1):347. doi: 10.1186/s12913-023-09131-0. PMID: 36822645; PMCID: PMC9932380.
28. De Almeida LM, Almeida PC, Santos CB. A review of social determinants of health inequities in Brazil. Cien Saude Colet. 2023 Apr;28(4):1461-1475. doi: 10.1590/1413-81232023284.07512023. PMID: 36985201.
29. Rahman SM, Ahinkorah BO, Adu C. Socioeconomic determinants of maternal health care utilization in Ghana: a systematic review and meta-analysis. BMC Health Serv Res. 2023 Aug 1;23(1):818. doi: 10.1186/s12913-023-08962-0. PMID: 37423973; PMCID: PMC10338961.
30. Zawiah M, Hamid K, Khatib N. Barriers to Healthcare Access among Rural Women in Southeast Asia: A Systematic Review. BMC Health Serv Res. 2023 Jul 14;23(1):539. doi: 10.1186/s12913-023-09087-4. PMID: 37420893; PMCID: PMC10356529.
31. Ismail SA, Zodpey SP, Mohamed J. Exploring the Social Determinants of Maternal Health and Its Implications in India: A Systematic Review. J Community Health. 2024 Apr;49(2):340-355. doi: 10.1007/s10900-023-01431-7. Epub 2023 Oct 7. PMID: 37841743.